

Chebyshev's bias for composite numbers with restricted prime divisors

Pieter Moree

Abstract

Let $\pi(x; d, a)$ denote the number of primes $p \leq x$ with $p \equiv a \pmod{d}$. Chebyshev's bias is the phenomenon that 'more often' $\pi(x; d, n) > \pi(x; d, r)$, than the other way around, where n is a quadratic non-residue mod d and r is a quadratic residue mod d . If $\pi(x; d, n) \geq \pi(x; d, r)$ for every x up to some large number, then one expects that $N(x; d, n) \geq N(x; d, r)$ for every x . Here $N(x; d, a)$ denotes the number of integers $n \leq x$ such that every prime divisor p of n satisfies $p \equiv a \pmod{d}$. In this paper we develop some tools to deal with this type of problem and apply them to show that, for example, $N(x; 4, 3) \geq N(x; 4, 1)$ for every x . In the process we express the so called second order Landau-Ramanujan constant as an infinite series and show that the same type of formula holds true for a much larger class of constants.

Keywords: Comparative number theory, constants, primes in progression, multiplicative functions.

1 Introduction

Especially for small moduli d primes seem to have a preference for non-quadratic residue classes mod d over quadratic residue classes mod d . This phenomenon is called Chebyshev's bias [2]. For example, $\pi(x; 3, 1)$ does not exceed $\pi(x; 3, 2)$ for the first time until $x = 608981813029$, as was shown by Bays and Hudson [1]. On the other hand Littlewood [15] has shown that the function $\pi(x; 3, 2) - \pi(x; 3, 1)$ has infinitely many sign changes. Rubinstein and Sarnak [28] quantified some biases under the assumption of the Generalized Riemann Hypothesis (GRH) and the assumption that the non-negative imaginary parts of the nontrivial zeros of all Dirichlet L-functions are linearly independent over the rationals. Define δ_{q, a_1, a_2} to be the logarithmic density of the set of real numbers x such that the inequality $\pi(x; q, a_1) > \pi(x; q, a_2)$ holds, where the logarithmic density of a set S is

$$\lim_{x \rightarrow \infty} \frac{1}{\log x} \int_{[2, x] \cap S} \frac{dt}{t},$$

P. Moree: KdV Institute, University of Amsterdam, Plantage Muidersgracht 24, 1018 TV Amsterdam, The Netherlands, e-mail: moree@science.uva.nl

Mathematics Subject Classification (2000). 11N37, 11Y60, 11N13

assuming the limit exists. Under the aforementioned assumptions and assuming that $(\mathbb{Z}/q\mathbb{Z})^*$ is cyclic, Rubinstein and Sarnak showed that δ_{q,a_1,a_2} always exists and is strictly positive and, moreover, that $\delta_{q,n,r} > 0.5$ if and only if n is a non-square mod q and r is a square mod q . They calculated, amongst others, that $\delta_{4,3,1} = 0.9959 \dots$ and $\delta_{3,2,1} = 0.9990 \dots$. Thus Chebyshev's bias is not only an initial interval phenomenon. The comparison of the behaviour of primes lying in various arithmetic progressions is the subject of comparative prime number theory, which was systematically developed in a series of papers by Knapowski and Turán, cf. [34, 35]. More recent references are e.g. [16, 28] and various papers of J. Kaczorowski. One of the older papers, by Wintner [37], was inspired, interestingly enough, on the observation (p. 240) that there is 'an apparent parrallelism between certain problems in celestial mechanics on the one hand and "wobbly" terms of the explicit formula of Riemann and Von Mangoldt on the other hand'. Wintner, who has written many papers on celestial mechanics then could put his expertise in that field to good use.

Put $g_{d,a}(n) = 0$ if n has no prime divisor p satisfying $p \equiv a \pmod{d}$ and $g_{d,a}(n) = 1$ otherwise (note that $g_{d,a}(1) = 1$). We let $N(x; d, a) = \sum_{n \leq x} g_{d,a}(n)$. The contribution of the small primes to the growth of $N(x; d, a)$ is much bigger than to $\pi(x; d, a)$ and hence we might expect that if $\pi(x; d, a) \geq \pi(x; d, b)$ up to some reasonable x , then actually $N(x; d, a) \geq N(x; d, b)$ for every x . In general, given two non-negative multiplicative functions f and g such that f and g are equal to a positive constant τ in the primes on average and such that there is a bias towards f in the sense that $\sum_{p \leq x} f(p) \geq \sum_{p \leq x} g(p)$ for all x up to some rather large number, is it true that $\sum_{n \leq x} f(n) \geq \sum_{n \leq x} g(n)$ for *every* x ? The asymptotic behaviour of the latter type of sums is well-understood and so proving that these types of results are true *asymptotically* is usually not difficult. We can for example invoke the following classical result due to Wirsing [38].

Theorem 1 (Wirsing [38]). *Let f be a multiplicative function satisfying $0 \leq f(p^r) \leq c_1 c_2^r$, $c_1 \geq 1$, $1 \leq c_2 < 2$, and $\sum_{p \leq x} f(p) = (\tau + o(1))x / \log x$, where τ, c_1 and c_2 are constants. Then, as $x \rightarrow \infty$,*

$$\sum_{n \leq x} f(n) \sim \frac{e^{-\gamma\tau}}{\Gamma(\tau)} \frac{x}{\log x} \prod_{p \leq x} \left(1 + \frac{f(p)}{p} + \frac{f(p^2)}{p^2} + \frac{f(p^3)}{p^3} + \dots \right),$$

where γ is Euler's constant and $\Gamma(\tau)$ denotes the gamma-function. (Here and in the sequel the letter p is used to indicate primes.)

We thus see that, for $i = 1$ and $i = 2$,

$$N(x; 3, i) \sim \frac{e^{-\gamma/2}}{\sqrt{\pi}} \frac{x}{\log x} \prod_{\substack{p \leq x \\ p \equiv i \pmod{3}}} \left(1 - \frac{1}{p} \right)^{-1}, \quad (1)$$

showing clearly the strong influence of the smaller primes. By [36, Theorem 2] we deduce from the latter formula that $N(x; 3, i) \sim C_{3,i} x / \sqrt{\log x}$, with

$$C_{3,1} = \frac{3^{1/4}}{\pi \sqrt{2}} \prod_{p \equiv 1 \pmod{3}} \left(1 - \frac{1}{p^2} \right)^{-1/2} = \frac{\sqrt{2}}{3^{5/4}} \prod_{p \equiv 2 \pmod{3}} \left(1 - \frac{1}{p^2} \right)^{\frac{1}{2}}, \quad (2)$$

where in the derivation of the (2) we used Euler's identity $\pi^2/6 = \prod_p (1 - p^{-2})^{-1}$ (another, selfcontained, derivation of (2) is given in Section 6). Using Merten's theorem or [36, Theorem 2] again, we easily infer that $C_{3,2} = 2/(3\pi C_{3,1})$. Restricting to the primes $p \leq 29$, we compute that $C_{3,1} < 0.302$ and $C_{3,2} > 0.703$ (for more precise numerical evaluations see Section 6). We thus infer that $N(x; 3, 2) \geq N(x; 3, 1)$ for every sufficiently large x . If we want to make this effective, the extensive literature, cf. [24], on multiplicative functions satisfying conditions as in Wirsing's theorem seems to offer no help as nobody seems to have been concerned with proving effective results in this area, which is precisely what the Chebyshev bias problem for composites challenges us to do. In this paper we develop some tools for this and apply them to prove:

Theorem 2 *The inequalities $N(x; 3, 2) \geq N(x; 3, 1)$, $N(x; 4, 3) \geq N(x; 3, 1)$, $N(x; 3, 2) \geq N(x; 4, 1)$ and $N(x; 4, 3) \geq N(x; 4, 1)$ hold true for every x .*

Not surprisingly the Chebyshev bias problem for composites is rather computational in nature and this appears to preclude one from proving more general results.

The counting functions appearing in Theorem 2 can be shown to satisfy more precise asymptotic estimates than (1). Theorem 3 together with the prime number theorem for arithmetic progressions, shows that there exist constants $C_{d,a}, C_{d,a}(1), C_{d,a}(2), \dots$ such that for each integer $m \geq 0$ we have

$$N(x; d, a) = \frac{C_{d,a}x}{\sqrt{\log x}} \left(1 + \sum_{j=0}^m \frac{C_{d,a}(j)}{\log^j x} + O\left(\frac{1}{\log^{m+1} x}\right) \right),$$

where the implied constant may depend on m, a and d . Thus $N(x; d, a)$ satisfies an asymptotic expansion in the sense of Poincaré in terms of $\log x$. The most famous example of such an expression states that for $B(x)$, the counting function of the integers that can be represented as a sum of two integer squares, we have

$$B(x) = \frac{Kx}{\sqrt{\log x}} \left(1 + \sum_{j=0}^m \frac{K_{j+2}}{\log^j x} + O\left(\frac{1}{\log^{m+1} x}\right) \right), \quad (3)$$

where K is the Landau-Ramanujan constant and K_2 the second order Landau-Ramanujan constant. The Landau-Ramanujan constant is named after Landau [14] who proved in 1908, using contour integration, that $B(x) \sim Kx/\sqrt{\log x}$ and Ramanujan, who in his first letter to Hardy claimed he could prove that $B(x) = K \int_2^x dt/\sqrt{\log t} + O(x^{1/2+\epsilon})$, cf. [19]. Ramanujan's claim implies $K_2 = 1/2$ by partial integration, which was shown to be false by Shanks [32]. Indeed, we have

$$K = \frac{1}{\sqrt{2}} \prod_{p \equiv 3 \pmod{4}} \left(1 - \frac{1}{p^2} \right)^{-1/2} = 0.76422365358922066299069873125 \dots$$

and

$$K_2 = \frac{1}{2} - \frac{\gamma}{4} - \frac{L'(1, \chi_4)}{4L(1, \chi_4)} + \frac{\log 2}{4} + \frac{1}{2} \sum_{p \equiv 3 \pmod{4}} \frac{\log p}{p^2 - 1} = 0.5819486593172907 \dots$$

These constants have been calculated with 1000D precision at least, see [11]. It was a folklore result that $B(x)$ should satisfy (3), which was written down by Serre [31], who gave some nice applications to fourier coefficients of modular forms as well.

Let f be a non-negative multiplicative function. Suppose there exists a positive constant τ such that

$$\sum_{n \leq x} f(n) = \lambda_1(f) x \log^{\tau-1} x \left(1 + (1 + o(1)) \frac{\lambda_2(f)}{\log x} \right), \quad x \rightarrow \infty.$$

We then define $\lambda_2(f)$ to be the *generalized second-order Landau-Ramanujan constant*. In Theorem 4 we will identify a subclass of multiplicative functions for which this constant exists and express it as an infinite series.

The second-order generalized Landau-Ramanujan constant $\lambda_2(f)$ is closely related to the constant B_f appearing in the proof of Lemma 1 (the key lemma in the proof of Theorem 2). (I suggest to read the next section first before reading further.) Lemma 1 yields an effective estimate for $\mu_f(x)$, provided we can find constants τ, C_- and C_+ satisfying (7). For the functions f associated to the quantities in Theorem 2 we find admissible values of these constants in Section 8, which requires effective estimates for counting functions of squarefree numbers of a certain type (Section 7). (At the end of Section 8 we show that under GRH finding C_- and C_+ is much easier.) In Section 4 we show how to obtain effective estimates for $M_f(x)$ from effective estimates for $\mu_f(x)$. In Section 9 we show how to prove certain subcases of Theorem 2 for every x up to some large x_0 using existing numerical work on the associated Chebyshev prime biases. All these ingredients then come together in Section 10, where a proof of Theorem 2 is given.

In Section 5 we find an infinite series expansion for the constant B_f appearing in Lemma 2 (we have $C_- \leq B_f \leq C_+$) and relate it to the generalized second-order Landau-Ramanujan constant. Section 6 contains a numerical study of some of the constants appearing in this paper.

In [20] the methods developed in this paper are somewhat refined and then used to resolve Schmutz Schaller's conjecture (see [29, p. 201] or the introduction of [6]) that the hexagonal lattice is "better" than the square lattice. More precisely, let $0 < h_1 < h_2 < \dots$ be the positive integers, listed in ascending order, which can be written as $h_i = x^2 + 3y^2$ for integers x and y . Let $0 < q_1 < q_2 < \dots$ be the positive integers, listed in ascending order, which can be written as $q_i = x^2 + y^2$ for integers x and y . Then Schmutz Schaller's conjecture is that $q_i \leq h_i$ for $i = 1, 2, 3, \dots$.

2 Notation

Let f be a non-negative real-valued multiplicative function. We define $M_f(x) = \sum_{n \leq x} f(n)$, $\mu_f(x) = \sum_{n \leq x} f(n)/n$ and $\lambda_f(x) = \sum_{n \leq x} f(n) \log n$. We denote the formal Dirichlet series $\sum_{n=1}^{\infty} f(n)n^{-s}$ associated to f by $L_f(s)$. If $f(p)$ equals $\tau > 0$ on average at primes p , it can be shown that $\lim_{s \rightarrow 1+0} (s-1)^{\tau} L_f(s)$ exists, under some mild additional conditions on f . In that case we put

$$C_f := \frac{1}{\Gamma(\tau)} \lim_{s \rightarrow 1+0} (s-1)^{\tau} L_f(s).$$

We have $C_f > 0$. We define $\Lambda_f(n)$ by

$$-\frac{L'_f(s)}{L_f(s)} = \sum_{n=1}^{\infty} \frac{\Lambda_f(n)}{n^s}.$$

Notice that

$$f(n) \log n = \sum_{d|n} f(d) \Lambda_f\left(\frac{n}{d}\right). \quad (4)$$

The notation suggests that $\Lambda_f(n)$ is an analogue of the Von Mangoldt function. Indeed, if $f = \mathbf{1}$, then $L_f(s) = \zeta(s)$ and $\Lambda_f(n) = \Lambda(n)$. From (4) we infer by Möbius inversion the well-known formula

$$\Lambda(n) = \sum_{d|n} \mu(d) \log \frac{n}{d}. \quad (5)$$

In general, on writing $L_f(s)$ as an Euler product, one easily sees that $\Lambda_f(n)$ is zero if n is not a prime power. If f is the characteristic function of a subsemigroup of the natural integers with $(1 <) q_1 < q_2 < \dots$ as generators, then it can be shown that $\Lambda_f(n) = \log q_i$ if n equals a positive power of a generator q_i and $\Lambda_f(n) = 0$ otherwise. Thus for example, if $f = g_{d,a}$, then $\Lambda_{g_{d,a}}(n) = \log p$ if $n = p^r$, $r \geq 1$ and $p \equiv a \pmod{d}$, and $\Lambda_{g_{d,a}}(n) = 0$ otherwise.

From property (4) of $\Lambda_f(n)$, we easily infer that

$$\lambda_f(x) = \sum_{n \leq x} f(n) \psi_f\left(\frac{x}{n}\right), \quad (6)$$

where $\psi_f(x) = \sum_{n \leq x} \Lambda_f(n)$. For some further properties of $\Lambda_f(n)$ the reader is referred to [19, §2.2].

The notation x_0, α and β is used to indicate inessential local constants, their values might be different in different contexts.

3 Effective estimates for $\mu_f(x)$

The following result will play a crucial rôle. It uses some ideas from the proof of Theorem A in [33].

Lemma 1 *Let f be a non-negative multiplicative arithmetic function. Suppose that there exists constants $\tau(> 0)$, C_- and C_+ such that*

$$C_- \leq \sum_{n \leq x} \frac{\Lambda_f(n)}{n} - \tau \log x \leq C_+ \quad \text{for every } x \geq 1, \quad (7)$$

then, for $x > \exp(C_+)$, we have

$$\frac{C_f}{\tau} \log^\tau x \frac{\left(1 - \frac{C_+}{\log x}\right)^{\tau+1}}{1 - \frac{C_-}{\log x}} \leq \mu_f(x) \leq \frac{C_f}{\tau} \log^\tau x \frac{\left(1 - \frac{C_-}{\log x}\right)^{\tau+1}}{1 - \frac{C_+}{\log x}} \quad (8)$$

where

$$C_f := \frac{1}{\Gamma(\tau)} \lim_{s \rightarrow 1+0} (s-1)^\tau L_f(s). \quad (9)$$

Remark. An alternative expression for C_f is given by

$$C_f = \frac{1}{\Gamma(\tau)} \lim_{s \rightarrow 1+0} \prod_p \left(1 + \sum_{k=1}^{\infty} \frac{f(p^k)}{p^{ks}} \right) \left(1 - \frac{1}{p^s} \right)^{\tau}.$$

Proof of Lemma 1. Let B_f be an arbitrary constant and write

$$\sum_{n \leq x} \frac{\Lambda_f(n)}{n} = \tau \log x + B_f + E_f(x). \quad (10)$$

(This is unnecessary for this proof, but needed in the proof of Lemma 3, so we do this now to save some space later.) We have

$$\begin{aligned} \sum_{n \leq x} \frac{f(n) \log n}{n} &= \sum_{n \leq x} \sum_{d|n} \frac{f(d)}{d} \frac{\Lambda_f(n/d)}{\frac{n}{d}} = \sum_{d \leq x} \frac{f(d)}{d} \sum_{k \leq \frac{x}{d}} \frac{\Lambda_f(k)}{k} \\ &= \tau \sum_{n \leq x} \frac{f(n)}{n} \log \left(\frac{x}{n} \right) + B_f \mu_f(x) + \sum_{n \leq x} \frac{f(n)}{n} E_f \left(\frac{x}{n} \right). \end{aligned}$$

We write this equality in the form

$$-\sum_{n \leq x} \frac{f(n)}{n} \log \left(\frac{x}{n} \right) + \mu_f(x) \log x = \tau \sum_{n \leq x} \frac{f(n)}{n} \log \left(\frac{x}{n} \right) + B_f \mu_f(x) + \sum_{n \leq x} \frac{f(n)}{n} E_f \left(\frac{x}{n} \right).$$

This inequality on its turn can be written, using that

$$\sum_{n \leq x} \frac{f(n)}{n} \log \frac{x}{n} = \sum_{n \leq x} \frac{f(n)}{n} \int_n^x \frac{dt}{t} = \int_1^x \frac{\mu_f(t)}{t} dt,$$

as

$$\mu_f(x) \log x - (\tau + 1) \int_1^x \frac{\mu_f(v)}{v} dv = B_f \mu_f(x) + \sum_{n \leq x} \frac{f(n)}{n} E_f \left(\frac{x}{n} \right). \quad (11)$$

Put $\sigma_f(x) = \int_1^x \frac{\mu_f(v)}{v} dv$. By assumption $C_- \leq B_f + E_f(x) \leq C_+$ for $x \geq 1$. Using (11) we then deduce that $\mu_f(x) = (\tau + 1)\sigma_f(x)/\log x + \mu_f(x)\epsilon_f(x)$, where $C_- \leq \epsilon_f(x) \log x \leq C_+$. Solving this for $\mu_f(x)$ we find that

$$\mu_f(x) = \frac{1}{1 - \epsilon_f(x)} \frac{\tau + 1}{\log x} \sigma_f(x), \quad x \geq x_0, \quad (12)$$

where $x_0 := \exp((1 + \delta)C_+)$, and $\delta > 0$ is arbitrary and fixed. In the rest of the proof we assume that $x \geq x_0$. Let

$$R_f(t) := \log \left(\frac{\tau + 1}{\log^{\tau+1} t} \sigma_f(t) \right).$$

Note that, for $t \geq x_0$,

$$R'_f(t) = \frac{\tau + 1}{t \log t} \frac{\epsilon_f(t)}{[1 - \epsilon_f(t)]} \quad (13)$$

and hence $R'_f(t) = O(t^{-1} \log^{-2} t)$. Thus $\int_x^\infty R'_f(t) dt$ converges absolutely, and therefore $\int_x^\infty R'_f(t) dt = A_f - R_f(x)$, for some constant A_f not depending on x . On writing $D_f = \exp(A_f)$ we obtain

$$\frac{\tau+1}{\log^{\tau+1} x} \sigma_f(x) = \exp(R_f(x)) = D_f \exp\left(-\int_x^\infty R'_f(t) dt\right) \quad (14)$$

Using (13) and $C_- \leq \epsilon_f(x) \log x \leq C_+$, we see that

$$\int_x^\infty \frac{C_-(\tau+1)}{t \log t [\log t - C_-]} dt \leq \int_x^\infty R'_f(t) dt \leq \int_x^\infty \frac{C_+(\tau+1)}{t \log t [\log t - C_+]} dt.$$

Thus

$$-(\tau+1) \log\left(1 - \frac{C_-}{\log x}\right) \leq \int_x^\infty R'_f(t) dt \leq -(\tau+1) \log\left(1 - \frac{C_+}{\log x}\right).$$

On combining (14) with (13) we deduce that

$$D_f \left(1 - \frac{C_+}{\log x}\right)^{\tau+1} \leq \frac{\tau+1}{\log^{1+\tau} x} \sigma_f(x) \leq D_f \left(1 - \frac{C_-}{\log x}\right)^{\tau+1} \quad (15)$$

We will now show that $D_f = C_f/\tau$. The inequalities (15) in combination with (12) imply in particular that

$$\mu_f(x) = D_f \log^\tau x + O(\log^{\tau-1} x). \quad (16)$$

By partial integration and using the well-known integral expression for the gamma function we find that

$$\begin{aligned} L_f(s) &= (s-1) \int_1^\infty \frac{\mu_f(t)}{t^s} dt = (s-1) \int_1^\infty \frac{D_f \log^\tau t + O(\log^{\tau-1} t)}{t^s} dt \\ &= D_f \frac{\Gamma(\tau+1)}{(s-1)^\tau} + O\left(\frac{s-1}{(s-1)^\tau}\right) \end{aligned}$$

and thus $D_f = C_f/\tau$. The inequalities (15) together with (12) yield (8) on using that $D_f = C_f/\tau$ and $C_- \leq \epsilon_f(x) \log x \leq C_+$. $\square$

The convolutional nature of $\sum_{n \leq x} E(x/n) f(n)/n$ forces us to require that $x \geq 1$ in (7) (whereas we would like to replace it with $x \geq x_0$). Nevertheless we can invoke the following easy lemma to improve on (8).

Lemma 2 *Suppose that there exists constants D_- and D_+ such that for every $x \geq x_0$,*

$$D_- \mu_f(x) \leq B_f \mu_f(x) + \sum_{n \leq x} \frac{f(n)}{n} E_f\left(\frac{x}{n}\right) \leq D_+ \mu_f(x). \quad (17)$$

Then we have, for $x > \max\{x_0, \exp(D_+)\}$,

$$\frac{C_f}{\tau} \log^\tau x \frac{\left(1 - \frac{D_+}{\log x}\right)^{\tau+1}}{1 - \frac{D_-}{\log x}} \leq \mu_f(x) \leq \frac{C_f}{\tau} \log^\tau x \frac{\left(1 - \frac{D_-}{\log x}\right)^{\tau+1}}{1 - \frac{D_+}{\log x}}. \quad (18)$$

Proof. Follows easily on closer scrutiny of the previous proof. $\square$

We now give an example of how Lemma 2 can be used. By assumption we have $E_f(x) \leq C_+ - B_f$ for every $x \geq 1$. Suppose that $E_f(x) \leq C'_+ - B_f$ for $x \geq n_0$, where $C'_+ < C_+$. An upper bound for the innerterm in (17) is then given by

$$C_+ \mu_f(x) - (C_+ - C'_+) \mu_f\left(\frac{x}{n_0}\right).$$

Using the explicit bounds in (8) we can then find an x_0 and $D_+ < C_+$ such that the conditions of Lemma 2 are satisfied (note that $D_+ > C'_+$). By applying (18) instead of (8) a better value for D_+ can then be obtained. Then iterate.

By making an assumption on $E_f(x)$ we will, not surprisingly, be able to do better than both Lemma 1 and Lemma 2.

Lemma 3 *Let f be a non-negative multiplicative arithmetic function and suppose that (10) holds with*

$$|E_f(x)| \leq \frac{c_0}{\max\{1, \log x\}}. \quad (19)$$

for every $x \geq 1$, where c_0 is some explicit constant. Then there exist effectively computable constants α, β and x_0 such that

$$\mu_f(x) = \frac{C_f}{\tau} \log^\tau x - C_f B_f \log^{\tau-1} x + \mathcal{E}_f(x),$$

where $\alpha \log^{\tau-1/2} x \leq \mathcal{E}_f(x) \leq \beta \log^{\tau-1/2} x$ for every $x \geq x_0$.

Proof. We denote the right hand side of (19) by $h(x)$ and put $s(x) = x/e^{\sqrt{\log x}}$. Let $x_0 \geq e$. Using Lemma 1 it is not difficult to see that

$$\beta_f(x_0) := \sup_{x \geq x_0} \sqrt{\log x} \left\{ 1 - \frac{\mu_f(s(x))}{\mu_f(x)} \left(1 - \frac{1}{\sqrt{\log x}} \right) \right\},$$

is finite and can be effectively computed (note that $\beta_f(x_0) \geq \tau + 1$).

Clearly

$$\left| \sum_{n \leq x} \frac{f(n)}{n} E_f\left(\frac{x}{n}\right) \right| \leq \sum_{n \leq s(x)} \frac{f(n)}{n} h\left(\frac{x}{n}\right) + \sum_{s(x) < n \leq x} \frac{f(n)}{n} h\left(\frac{x}{n}\right).$$

Denote the latter two sums by I_1 and I_2 . We have $I_1 \leq c_0 \mu_f(s(x))/\sqrt{\log x}$ and $I_2 \leq c_0(\mu_f(x) - \mu_f(s(x)))$. We thus find that (12) holds true with

$$|\epsilon_f(x) - \frac{B_f}{\log x}| \leq \frac{c_0 \beta_f(x_0)}{\log^{3/2} x}, \quad x \geq x_0.$$

Proceeding as in the proof of Lemma 1, but with this improved error estimate, the result then easily follows. $\square$

4 Relating $\mu_f(x)$ to $M_f(x)$

Given an effective estimate for $\mu_f(t)$, we can derive an effective estimate for $M_f(t)$ on using that

$$M_f(x) - M_f(x_0) = \int_{x_0}^x t \, d\mu_f(t). \quad (20)$$

Suppose that

$$\frac{C_f}{\tau} \log^\tau x \left(1 + \frac{\alpha}{\log x}\right) \leq \mu_f(x) \leq \frac{C_f}{\tau} \log^\tau x \left(1 + \frac{\alpha + \beta}{\log x}\right),$$

for some constants α and β and every $x \geq x_0$ (if the conditions of Lemma 1 are satisfied, such α, β and x_0 can certainly be determined). This leads to an upperbound for $M_f(x)$ that is asymptotically equal to $C_f(1 + \beta/\tau)x \log^{\tau-1} x$ and a lowerbound that is asymptotically equal to $\max\{0, C_f(1 - \beta/\tau)x \log^{\tau-1} x\}$. These estimates are too weak for our purposes.

Write $\mu_f(x) = C_f \log^\tau x / \tau - C_f B_f \log^{\tau-1} x + \mathcal{E}_f(x)$, cf. Lemma 3, and suppose that $\mathcal{E}_f^-(x) \leq \mathcal{E}_f(x) \leq \mathcal{E}_f^+(x)$ for every $x \geq x_0$, where $\mathcal{E}_f^+(x)$ and $\mathcal{E}_f^-(x)$ are effectively computable. (This supposition is certainly true if the conditions of Lemma 3 are satisfied.) Let $C_f(x_0) = M_f(x_0) - x_0 \mathcal{E}_f(x_0) - C_f x_0 \log^\tau x_0$. Then an easy computation shows that for every $x \geq x_0$,

$$M_f(x) \leq C_f x \log^{\tau-1} x + (1 - \tau)C_f(1 + B_f) \int_{x_0}^x \log^{\tau-2} t \, dt + C_f(x_0) + R_f(x),$$

where

$$x \mathcal{E}_f^-(x) - \int_{x_0}^x \mathcal{E}_f^+(t) \, dt \leq R_f(t) \leq x \mathcal{E}_f^+(x) - \int_{x_0}^x \mathcal{E}_f^-(t) \, dt.$$

There are various problems with this approach, one of the major ones being getting a good estimate for c_0 in Lemma 3.

An alternative approach starts with the observation that, for $x \geq 2$,

$$M_f(x) = \int_{2-}^x \frac{d\lambda_f(t)}{\log t} = \frac{\lambda_f(x)}{\log x} + \int_2^x \frac{\lambda_f(t)}{t \log^2 t} \, dt, \quad (21)$$

and that if we have explicit bounds of the type $\alpha x < \psi_f(x) < \beta x$, then $\lambda_f(x)$ can be related to $x\mu_f(x)$ by (6). Note in particular that if $\lambda_f(x) \geq \lambda_g(x)$ for every $x \geq 2$, then $M_f(x) \geq M_g(x)$ for every x (the reverse implication is not always true in general). The disadvantage of proving something stronger is hopefully compensated by the fact that $\lambda_f(x)$ can be easily related to $\mu_f(x)$.

5 The generalized second-order Ramanujan-Landau constant

In Theorem 4 we will identify a subclass of multiplicative functions for which the generalized Landau-Ramanujan constant (defined in Section 1) exists and relate it to an infinite series involving $\Lambda_f(n)$. The following result will play an essential rôle in this.

Theorem 3 [19, Theorem 6]. *Let f be a multiplicative function satisfying*

$$0 \leq f(p^r) \leq c_1 c_2^r, \quad c_1 \geq 1, \quad 1 \leq c_2 < 2, \quad (22)$$

and $\sum_{p \leq x} f(p) = \tau \text{Li}(x) + O(x \log^{-2-\rho} x)$, where τ and ρ are positive real fixed numbers. Then there exists a constant B_f such that (10) holds with $E_f(x) = O(\log^{-\rho} x)$. Moreover, for every $\epsilon > 0$,

$$\sum_{n \leq x} \frac{f(n)}{n} = \sum_{0 \leq \nu < \rho+1} a_\nu \log^{\tau-\nu} x + O(\log^{\tau-1-\rho+\epsilon} x), \quad (23)$$

where the implied constant depends at most on f and ϵ . In case f is completely multiplicative, condition (22) can be weakened to $\sum_{p, r \geq 2, p^r > x} (f(p)/p)^r \log p = O(\log^{-\rho} x)$.

Proof. This result is just Theorem 6 of [19], except for the claim regarding $E_f(x)$, the truth of which is however established in the course of the proof of Theorem 6 of [19]. $\square$

The next result shows that the second-order Landau-Ramanujan constant is closely related to the constant B_f appearing in (10).

Theorem 4 *Let f be a multiplicative function satisfying the hypothesis of Theorem 3 with $\rho > 1$. Then $\lambda_2(f)$, the generalized second-order Landau-Ramanujan constant, equals*

$$\lambda_2(f) = (1 - \tau) \left(1 + \tau\gamma + \sum_{n=1}^{\infty} \frac{\Lambda_f(n) - \tau}{n} \right),$$

or alternatively $\lambda_2(f) = (1 - \tau)(1 + B_f)$, where

$$B_f := \lim_{x \rightarrow \infty} \left(\sum_{n \leq x} \frac{\Lambda_f(n)}{n} - \tau \log x \right).$$

Proof. Since by assumption $\rho > 1$, we have by (23)

$$\mu_f(x) = a_0 \log^\tau x + a_1 \log^{\tau-1} x + a_2 \log^{\tau-2} x + O(\log^{\tau-2-\delta} x),$$

for some $\delta > 0$. Theorem 3 implies that B_f exists. Using that $\log x = \sum_{n \leq x} 1/n - \gamma + o(1)$, we see that it suffices to prove that $\lambda_2(f) = (1 - \tau)(1 + B_f)$. Theorem 3 yields that $E_f(x) = O(\log^{-1} x)$, hence the conditions of Lemma 3 are satisfied and it follows that $a_0 = \tau C_f$ and $a_1 = -C_f B_f$. On using that $M_f(x) = x\mu_f(x) - \int_1^x \mu_f(t) dt$ it follows by partial integration that $\lambda_1(f) = C_f$ and $\lambda_2(f) = (1 - \tau)(1 + B_f)$, as required. $\square$

Example. Let b_1 be the characteristic function of the set of natural numbers that can be written as a sum of two integer squares. This is a subsemigroup of the natural numbers that is generated by the primes p with $p \equiv 1 \pmod{4}$, $p = 2$ and

the squares of the remaining prime numbers (this result goes back to Fermat). By what has been said in Section 2, it then follows that

$$\Lambda_{b_1}(n) = \begin{cases} 2 \log p & \text{if } n = p^r, \ r \geq 1 \text{ and } p \equiv 3 \pmod{4}; \\ \log p & \text{if } n = p^r, \ r \geq 1 \text{ and } p \equiv 1 \pmod{4} \text{ or } p = 2; \\ 0 & \text{otherwise.} \end{cases}$$

Application of Theorem 4 yields the following two formulae for the second-order Landau-Ramanujan constant K_2 (cf. Section 1):

$$K_2 = \frac{1}{2} \left(1 + \frac{\gamma}{2} + \sum_{n=1}^{\infty} \frac{\Lambda_{b_1}(n) - \frac{1}{2}}{n} \right) = \frac{1}{2} \lim_{x \rightarrow \infty} \left(1 + \sum_{n \leq x} \frac{\Lambda_{b_1}(n)}{n} - \frac{1}{2} \log x \right).$$

6 Numerical evaluation of certain constants

In order to complete our proof we need to evaluate certain constants with enough precision. For some of them this has been done before.

We first consider the evaluation of $C_{3,1}$ and $C_{3,2}$. We have, for $\Re(s) > 1$, $L_{g_{3,1}}(s) = \prod_{p \equiv 1 \pmod{3}} (1 - p^{-s})^{-1}$. Note that

$$L_{g_{3,1}}(s)^2 = \zeta(s) L(s, \chi_3) (1 - 3^{-s}) \prod_{p \equiv 2 \pmod{3}} (1 - p^{-2s}). \quad (24)$$

From this, (9), $\lim_{s \rightarrow 1+0} (s-1)\zeta(s) = 1$ and the fact that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$, we obtain

$$C_{3,1}^2 = \frac{2L(1, \chi_3)}{3\pi} \prod_{p \equiv 2 \pmod{3}} \left(1 - \frac{1}{p^2} \right).$$

If χ is a real primitive character modulo k and $\chi(-1) = -1$, then

$$L(1, \chi) = -\frac{\pi}{k^{3/2}} \sum_{n=1}^k n \chi(n),$$

by Dirichlet's celebrated class number formula (cf. equation (17) of Chapter 6 of [8]). We infer that $L(1, \chi_3) = \pi/\sqrt{27}$. Using that $C_{g_{3,1}} \geq 0$ and $\zeta(2) = \pi^2/6$, we then deduce (2). Using that $L_{g_{3,2}}(s) L_{g_{3,1}}(s) (1 - 3^{-s})^{-1} = \zeta(s)$, we infer that $C_{3,2} = 2/(3\pi C_{3,1})$.

In order to compute $C_{3,2}$ and $C_{3,1}$ with many decimal accuracy we proceed as in Shanks [32, p. 78]. We note that, for $\Re(s) > 1/2$,

$$\prod_{p \equiv 2 \pmod{3}} (1 - p^{-2s})^2 = \frac{L(2s, \chi_3)}{\zeta(2s)(1 - 3^{-2s})} \prod_{p \equiv 2 \pmod{3}} (1 - p^{-4s}), \quad (25)$$

from which we infer by recursion that

$$C_{3,1} = \frac{\sqrt{2}}{3^{\frac{5}{4}}} \prod_{n=1}^{\infty} \left(\frac{L(2^n, \chi_3)}{(1 - 3^{-2^n}) \zeta(2^n)} \right)^{\frac{1}{2^{n+1}}}.$$

Because of the lacunary character of this expression, it can be calculated quickly up to high precision, which yields $C_{3,1} = 0.3012165544749342124 \dots$ and $C_{3,2} = 0.7044984335 \dots$. Similarly one can show that $C_{4,3} = 1/(2\pi C_{4,1})$ and

$$C_{4,1} = \frac{1}{2\sqrt{2}} \prod_{p \equiv 3 \pmod{4}} \left(1 - \frac{1}{p^2}\right)^{1/2} = \frac{1}{\pi} \prod_{p \equiv 1 \pmod{4}} \left(1 - \frac{1}{p^2}\right)^{-1/2}.$$

Using Shanks' trick we then infer that $C_{4,1} = 0.3271293669410263824002328 \dots$ and $C_{4,3} = 0.4865198883 \dots$.

On noting that, for $\Re(s) \geq 1$,

$$\sum_{n=1}^{\infty} \frac{\Lambda(n) - 1}{n^s} = -\frac{\zeta'(s)}{\zeta(s)} - \zeta(s),$$

and using that $\zeta(s) = 1/(s-1) + \gamma + O(s-1)$ is the Taylor series for $\zeta(s)$ around $s = 1$ (see e.g. [22, pp. 162-164]), one infers that

$$\sum_{n \leq x} \frac{\Lambda(n)}{n} = \sum_{n \leq x} \frac{1}{n} - 2\gamma + o(1) = \log x - \gamma + o(1). \quad (26)$$

Taking the logarithmic derivative of (24) one obtains that

$$-2 \frac{L'_{g_{3,1}}(s)}{L_{g_{3,1}}(s)} = -\frac{\zeta'(s)}{\zeta(s)} - \frac{L'}{L}(s, \chi_3) - \frac{\log 3}{3^s - 1} - 2 \sum_{p \equiv 2 \pmod{3}} \frac{\log p}{p^{2s} - 1},$$

from which one easily infers that

$$2 \sum_{n \leq x} \frac{\Lambda_{g_{3,1}}(n)}{n} = \sum_{n \leq x} \frac{\Lambda(n)}{n} - \frac{L'}{L}(1, \chi_3) - \frac{\log 3}{2} - 2 \sum_{p \equiv 2 \pmod{3}} \frac{\log p}{p^2 - 1} + o(1),$$

which yields, on invoking (26),

$$2B_{g_{3,1}} = -\gamma - \frac{L'}{L}(1, \chi_3) - \frac{\log 3}{2} - 2 \sum_{p \equiv 2 \pmod{3}} \frac{\log p}{p^2 - 1}.$$

Similarly we deduce that

$$2B_{g_{4,1}} = -\gamma - \frac{L'}{L}(1, \chi_4) - \log 2 - 2 \sum_{p \equiv 3 \pmod{4}} \frac{\log p}{p^2 - 1}.$$

As to the numerical evaluation of $B_{g_{3,1}}$ and $B_{g_{4,1}}$, we note that

$$2 \sum_{p \equiv 2 \pmod{3}} \frac{\log p}{p^2 - 1} = -\frac{d}{ds} \log \prod_{p \equiv 2 \pmod{3}} \left(\frac{1}{1 - p^{-2s}} \right) \Big|_{s=1}.$$

Then, applying (25) m times, we obtain

$$\sum_{p \equiv 2 \pmod{3}} \frac{\log p}{p^2 - 1} = \sum_{p \equiv 2 \pmod{3}} \frac{\log p}{p^{2^{m+1}} - 1} + \frac{1}{2} \sum_{n=1}^m \left\{ \frac{L'}{L}(2^n, \chi_3) - \frac{\zeta'}{\zeta}(2^n) - \frac{\log 3}{3^{2^n} - 1} \right\}.$$

Now L -functions and their derivatives can be computed with high accuracy using for example PARI (cf. [3, Section 10.3]). On doing so we find that the prime sum in the left hand side of the latter formula equals $0.3516478132638087560157790\dots$. Similarly we have

$$\sum_{p \equiv 3 \pmod{4}} \frac{\log p}{p^2 - 1} = \sum_{p \equiv 3 \pmod{4}} \frac{\log p}{p^{2^{m+1}} - 1} + \frac{1}{2} \sum_{n=1}^m \left\{ \frac{L'}{L}(2^m, \chi_4) - \frac{\zeta'}{\zeta}(2^m) - \frac{\log 2}{2^{2^m} - 1} \right\}.$$

We thus find that the sum on the left hand side equals $0.2287363531940324576\dots$. For more on evaluating infinite sums or products involving primes, we refer to [4] and [18].

For the logarithmic derivative $L'(1, \chi)/L(1, \chi)$ we find, with $\chi = \chi_3$ and $\chi = \chi_4$, $\frac{L'}{L}(1, \chi_3) = 0.36828161597014784263323790407578664254876430999\dots$ and

$$\frac{L'}{L}(1, \chi_4) = 0.2456095847773141723888166261790625184335337829549\dots$$

An alternative way of evaluating the latter two logarithmic derivatives is by relating them to the gamma function or the arithmetic-geometric-mean (AGM). We have (Berger (1883), Lerch (1897), de Séguier (1899) and Landau [13]),

$$\frac{L'}{L}(1, \chi_4) = \log \left(M(1, \sqrt{2})^2 \frac{e^\gamma}{2} \right),$$

where $M(1, \sqrt{2})$ denotes the limiting value of Lagrange's AGM algorithm $a_{n+1} = (a_n + b_n)/2$, $b_{n+1} = \sqrt{a_n b_n}$ with inputs $a_0 = 1$ and $b_0 = \sqrt{2}$. It can be shown that $M(1, \sqrt{2}) = \sqrt{\frac{2}{\pi}} \Gamma(\frac{3}{4})^2$. Gauss showed (in his diary), cf. [7], that

$$\frac{1}{M(1, \sqrt{2})} = \frac{2}{\pi} \int_0^1 \frac{dx}{\sqrt{1-x^4}}.$$

The total arclength of the lemniscate $r^2 = \cos(2\theta)$ is given by $2L$, where $L := \pi/M(1, \sqrt{2})$ is the so-called *lemniscate constant*. If $\chi = \chi_3$ we have similarly, with $z := \sin(\frac{\pi}{12}) = \frac{(\sqrt{3}-1)}{\sqrt{8}}$,

$$\frac{L'}{L}(1, \chi_3) = \log \left(\frac{2^{\frac{4}{3}} M(1+z, 1-z)^2 e^\gamma}{3} \right), \quad M(1+z, 1-z) = \frac{2^{\frac{4}{3}} \pi^2}{3^{\frac{1}{4}} \Gamma(\frac{1}{3})^3},$$

and

$$\frac{1}{M(1+z, 1-z)} = \frac{3^{\frac{1}{4}}}{\pi} \int_0^1 \frac{dx}{\sqrt{x(1-x^3)}}.$$

The values of $L'(1, \chi_4)$ and $L'(1, \chi_3)$ can also be determined using generalized Euler constants for arithmetical progressions, see Examples 1 and 2 of [9]. For general non-trivial real χ the quotients $L'(1, \chi)/L(1, \chi)$ 'feel' the zeroes of $L(s, \chi)$ close to 1 (see [8, pp. 80-83] for a quantitative version) and a study of their

average behaviour might throw some light on the (non)-existence of the Landau-Siegel zeros, cf. [17].

Putting our subcomputations together, we find that

$$B_{g_{3,1}} = -1.09904952586667653048446536830561 \dots$$

On noting that $\Lambda(n) = \Lambda_{g_{3,1}}(n) + \Lambda_{g_{3,2}}(n)$ if for $3 \nmid n$, it is easily deduced that $B_{g_{3,1}} = -\gamma - \frac{\log 3}{2} - B_{g_{3,2}}$, using which we compute that

$$B_{g_{3,2}} = -.02747228336891117581966934023805 \dots$$

Similarly, we find $B_{g_{4,1}} = -0.9867225683134286288516284 \dots$ and

$$B_{g_{4,3}} = -\log 2 - \gamma - B_{g_{4,1}} = -0.2836402771480495411721157 \dots$$

An alternative approach in calculating the constant $B_{g_{d,a}}$ is on invoking the formula

$$\varphi(d) \sum_{\substack{n \leq x \\ n \equiv a \pmod{d}}} \frac{\Lambda(n)}{n} = \log x - \gamma - \sum_{p|d} \frac{\log p}{p-1} - \sum_{\chi \neq \chi_0} \chi(\bar{a}) \frac{L'}{L}(1, \chi) + o(1),$$

where a and d are coprime integers, the sum is over the characters mod d different from the principal character and $\bar{a}$ is any integer such that $a\bar{a} \equiv 1 \pmod{d}$. The latter formula is derived by elementary means in [23].

Using Theorem 4 we are now in the position to compute some second-order Landau-Ramanujan constants. They are simply given by $\lambda_2(f) = (1 + B_f)/2$ for $f \in \{g_{3,1}, g_{3,2}, g_{4,1}, g_{4,3}\}$.

7 Effective estimates for squarefree integers

In the sequel we will establish some effective estimates for certain number theoretic functions of a real variable. The general procedure is to establish the estimates for every $x \geq x_0$ for some x_0 . The following lemma can then often be used to show that there exists a number $x_1 < x_0$ such that the estimates in fact hold true for every $x \geq x_1$. It reduces a seemingly continuous problem to a discrete one.

Lemma 4 *Let $y_1 > y_0$ be arbitrary real numbers. Let F and r be non-decreasing real-valued functions such that, moreover, F changes its value only at integers. Let $x_1, x_2, \dots, x_n$ be the integers in (y_0, y_1) where F changes its value. Put $x_0 = y_0$ and $x_{n+1} = y_1$. Then*

$$\sup_{y_0 \leq x \leq y_1} \{F(x) - r(x)\} = \max_{0 \leq i \leq n} \{F(x_i) - r(x_i)\}$$

and

$$\inf_{y_0 \leq x \leq y_1} \{F(x) - r(x)\} = \min_{0 \leq i \leq n} \{F(x_i) - r(x_{i+1})\}.$$

In our proof of Theorem 2 we need effective estimates for $Q_{\chi_3}(x)$ and $Q_{\chi_4}(x)$, where $Q_{\chi}(x)$ denotes the number of integers $n \leq x$ such that $\mu(n)\chi(n) \neq 0$. Note that Q_{χ_4} merely counts the odd squarefree numbers and hence we will use the more suggestive notation Q_{odd} for it. There are two obvious approaches in estimating these functions; relating them to $Q(x)$, where $Q(x)$ denotes the number of squarefree integers not exceeding x , and an ab initio approach. We demonstrate both approaches.

Put $R(x) = Q(x) - 6x/\pi^2$. It was shown by Moser and MacLeod [21] that $|R(x)| < \sqrt{x}$ for all x and that $|R(x)| < \sqrt{x}/2$ for $x \geq 8$. Cohen and Dress [5] showed that $|R(x)| < 0.1333\sqrt{x}$ for $x \geq 1664$.

Lemma 5 *For $x \geq 0$ we have*

$$\left| Q_{\chi_3}(x) - \frac{9}{2\pi^2}x \right| \leq 0.3154\sqrt{x} + 17.2,$$

and

$$\left| Q_{\chi_3}(x) - \frac{9}{2\pi^2}x \right| \leq \frac{1}{2}\sqrt{x} + 1. \quad (27)$$

Proof. We clearly have $Q(x) = Q_{\chi_3}(x) + Q_{\chi_3}(x/3)$, from which we infer that

$$Q_{\chi_3}(x) = \sum_{i=0}^{\infty} (-1)^i Q\left(\frac{x}{3^i}\right). \quad (28)$$

Put $x_0 = 1664$. On applying Lemma 4 with $y_0 = x_0/3$ and $y_1 = x_0$, we find that $|R(x)| \leq 0.15\sqrt{x}$ in the interval $(x_0/3, x_0]$. Similarly we compute that $|R(x)| \leq 0.29\sqrt{x}$ in the interval $(x_0/27, x_0/3]$. These estimates yield when combined with identity (28) and the quoted bounds for $|R(x)|$:

$$\begin{aligned} \left| Q_{\chi_3}(x) - \frac{9}{2\pi^2}x \right| &\leq \alpha \frac{\sqrt{3x}}{\sqrt{3}-1} + (0.15 - \alpha)\sqrt{x_0} + (0.29 - \alpha) \left(\sqrt{\frac{x_0}{3}} + \sqrt{\frac{x_0}{9}} \right) \\ &\quad + (0.5 - \alpha) \left(\sqrt{\frac{x_0}{27}} + \sqrt{\frac{x_0}{81}} \right) + (1 - \alpha)(\sqrt{9} + \sqrt{3} + 1 + \frac{1}{\sqrt{3}} + \frac{1}{3} + \cdots), \end{aligned}$$

where $\alpha = 0.1333$. The latter bound does not exceed $0.3154\sqrt{x} + 17.2$. From this bound we then infer that (27) holds for every $x \geq 10000$. We now apply Lemma 4 with $y_0 = 0$ and $y_1 = 10000$ to establish the validity of (27) in the remaining range. $\square$

Using that $|R(x)| \leq 0.15\sqrt{x}$ in the interval $(x_0/2, x_0]$ and $|R(x)| \leq 0.29\sqrt{x}$ in the interval $(x_0/32, x_0/2]$ we deduce, proceeding as in the proof of Lemma 5, that $|Q_{\text{odd}}(x) - 4x/\pi^2| \leq 0.4552\sqrt{x} + 26.5$. Although the latter bound is sharp enough for our purposes, we present a selfcontained proof of a slightly sharper bound (which uses ideas from [21]).

Lemma 6 *For $x \geq 0$ we have*

$$\left| Q_{\text{odd}}(x) - \frac{4}{\pi^2}x \right| \leq \frac{1}{2}\sqrt{x} + 1 \text{ and } \left| Q_{\text{odd}}(x) - \frac{4}{\pi^2}x \right| \leq \left(\frac{2}{\pi^2} + \frac{1}{4} \right)\sqrt{x} + \frac{1}{4}x^{\frac{1}{4}} + 2.$$

Proof. We have

$$Q_{\text{odd}}(x) = \sum_{\substack{n \leq x \\ n \text{ odd}}} |\mu(n)| = \sum_{\substack{n \leq x \\ n \text{ odd}}} \sum_{d^2 | n} \mu(d) = \sum_{\substack{d \leq x \\ d \text{ odd}}} \mu(d) \left[\frac{x}{2d^2} + \frac{1}{2} \right].$$

Put $R_{\text{odd}}(x) = Q_{\text{odd}}(x) - 4x/\pi^2$. On noting that $\sum_{d \text{ odd}} \mu(d)/d^2 = 8/\pi^2$, we find that

$$|R_{\text{odd}}(x)| \leq \left| \sum_{\substack{d^2 \leq x \\ d \text{ odd}}} \mu(d) \left(\frac{x}{2d^2} - \left[\frac{x}{2d^2} + \frac{1}{2} \right] \right) \right| + x \left| \sum_{\substack{d^2 > x \\ d \text{ odd}}} \frac{\mu(d)}{2d^2} \right|.$$

Since $|x - [x + 1/2]| \leq 1/2$ for every x , we deduce that

$$|R_{\text{odd}}(x)| \leq \frac{Q_{\text{odd}}(\sqrt{x})}{2} + x \left| \sum_{\substack{d^2 > x \\ d \text{ odd}}} \frac{\mu(d)}{2d^2} \right|. \quad (29)$$

Suppose that $x > 4$, then

$$\sum_{\substack{d^2 > x \\ d \text{ odd}}} \frac{1}{d^2} \leq \sum_{m > \frac{\sqrt{x}-1}{2}} \frac{1}{(2m+1)(2m-1)} = \sum_{m > \frac{\sqrt{x}-1}{2}} \left[\frac{1}{4m-2} - \frac{1}{4m+2} \right] \leq \frac{.5}{\sqrt{x}-2}.$$

On using this and the trivial estimate $Q_{\text{odd}}(x) \leq (x+1)/2$, we deduce that $|R_{\text{odd}}(x)| \leq \frac{1}{2}\sqrt{x} + 1$ on applying Lemma 4 with $y_0 = 0$ and $y_1 = 36$. Using the latter bound for $Q_{\text{odd}}(x)$ in (29) one then easily obtains the second stated bound in the formulation of the lemma on applying Lemma 4 with $y_0 = 0$ and $y_1 = 9$.

8 On the difference $\sum_{n \leq x} \frac{\Lambda_f(n)}{n} - \tau \log x$

In order to use Lemma 1 we need to find finite constants C_+ and C_- such that

$$C_- \leq \sum_{n \leq x} \frac{\Lambda_f(n)}{n} - \tau \log x \leq C_+$$

for every $x \geq 1$. Recall that $\psi_f(x) := \sum_{n \leq x} \Lambda_f(n)$. Suppose that $\psi_f(x) = \tau x + \mathcal{E}_f(x)$, where $|\mathcal{E}_f(x)| \leq c_\epsilon \log^{-1-\epsilon} x$ for $x \geq x_0$. Then

$$\sum_{n \leq x} \frac{\Lambda_f(n)}{n} = \tau \log x + B_f + \frac{\mathcal{E}_f(x)}{x} - \int_x^\infty \frac{\mathcal{E}_f(t)}{t^2} dt,$$

and thus, for $x \geq x_0$,

$$\left| \sum_{n \leq x} \frac{\Lambda_f(n)}{n} - \tau \log x - B_f \right| \leq \frac{c_\epsilon}{\log^\epsilon x} \left(\frac{1}{\epsilon} + \frac{1}{\log x} \right). \quad (30)$$

For example, if $f = \mathbf{1}$, it is known that $|\theta(x) - x| \leq 3.965x/\log^2 x$ for $x > 1$ [10, p.14]. Using this with the bound $\psi(x) - \theta(x) < 1.43\sqrt{x}$ [27, Theorem 13], we can compute C_+ and C_- in this case. Instead of carrying this out along these lines, we proceed slightly differently as this will result in a sharper bound for the difference in (30).

Lemma 7 For $x \geq 97$ we have

$$-\frac{1}{2\log x} + \frac{1}{2\sqrt{x}} \leq \sum_{n \leq x} \frac{\Lambda(n)}{n} - \log x + \gamma \leq \frac{2}{\sqrt{x}} + \frac{1}{2\log x}.$$

The upper bound holds even true for every $x > 1$.

Proof. By [27, Theorem 6] we have, for $x \geq 319$,

$$\left| \sum_{p \leq x} \frac{\log p}{p} - \log x - E \right| < \frac{1}{2\log x},$$

where $E = -\gamma - \sum_p \log p \sum_{k \geq 2} p^{-k}$. Notice that

$$\sum_{n \leq x} \frac{\Lambda(n)}{n} = \sum_{p \leq x} \frac{\log p}{p} + \sum_{k \geq 2} \sum_p \frac{\log p}{p^k} - \sum_{\substack{p^k > x \\ k \geq 2}} \frac{\log p}{p^k}.$$

By partial integration we find that

$$\sum_{\substack{p^k > x \\ k \geq 2}} \frac{\log p}{p^k} = \frac{\theta(x) - \psi(x)}{x} + \int_x^\infty \frac{\psi(t) - \theta(t)}{t^2} dt. \quad (31)$$

Suppose that $\alpha\sqrt{t} \leq \psi(t) - \theta(t) \leq \beta\sqrt{t}$ for $t \geq x_0$. Then, for $x \geq x_0$ the sum in (31) is in the interval $(\frac{2\alpha-\beta}{\sqrt{x}}, \frac{2\beta-\alpha}{\sqrt{x}})$. By Theorems 13 and 14 of [27] we can take $\alpha = 0.98$ and $\beta = 1.4262$ when $x_0 = 319$. On combining the various estimates, the result follows after some numerical analysis in the interval $(1, 319)$. $\square$

From Lemma 7 and Lemma 4 with $y_0 = 1$ and $y_1 = 215$ it is easily deduced that

$$\sup_{x \geq 1} \left\{ \sum_{n \leq x} \frac{\Lambda(n)}{n} - \log x \right\} = -\frac{\log 2}{2} = -0.34657359 \dots$$

and

$$\inf_{x \geq 1} \left\{ \sum_{n \leq x} \frac{\Lambda(n)}{n} - \log x \right\} = \frac{\log 2}{2} - \log 3 = -0.75203869 \dots$$

Other than for $f = \mathbf{1}$, the author is unaware of cases where an unconditional effective upper bound for $\mathcal{E}_f(x)$ of order $\log^{-1-\epsilon} x$ is known. Thus in order to obtain admissible values for C_+ and C_- in the case $f \in \{g_{3,1}, g_{3,2}, g_{4,1}, g_{4,3}\}$ we have to follow another approach, which is what we will do now. Notice that

$$2 \sum_{n \leq x} \frac{\Lambda_{g_{3,1}}(n)}{n} = \sum_{n \leq x} \frac{(1 + \chi_3(n))\Lambda(n)}{n} - 2 \sum_{\substack{p^r \leq \sqrt{x} \\ p \equiv 2 \pmod{3}}} \frac{\log p}{p^{2r}} - \sum_{1 < 3^r \leq x} \frac{\log 3}{3^r}. \quad (32)$$

The latter two sums are easily explicitly estimated and we already explicitly estimated $\sum_{n \leq x} \Lambda(n)/n$. If we can explicitly estimate $\sum_{n \leq x} \chi_3(n)\Lambda(n)/n$, we are done then. In order to achieve the latter goal, we need a few lemmas.

Lemma 8 Let h be a completely multiplicative function with $h(1) = 1$, then if $g(x) = \sum_{n \leq x} h(n)f(\frac{x}{n})$ for every x , it follows that $f(x) = \sum_{n \leq x} h(n)\mu(n)g(\frac{x}{n})$.

Proof. Substitute the expression $\sum_{mn \leq x} h(m)f(x/mn)$ for $g(x/n)$ in the sum $\sum_{n \leq x} h(n)\mu(n)g(x/n)$. The resulting expression simplifies to $f(x)$. $\square$

Lemma 9 Let χ be a non-principal character and $m_0 > 1$ be the smallest integer > 1 such that $\chi(m_0) \neq 0$. Then

$$\sum_{n \leq x} \frac{\chi(n)\Lambda(n)}{n} + \frac{L'}{L}(1, \chi) = O\left(\frac{L'}{L}(1, \chi) \frac{m_0}{x} Q_\chi\left(\frac{x}{m_0}\right)\right) + O\left(\frac{1}{x} \sum_{\substack{d \leq x/m_0 \\ \mu(d)\chi(d) \neq 0}} \log \frac{x}{d}\right).$$

Proof. On using (5) and writing $n = dd_1$ we obtain, for an arbitrary character χ ,

$$\sum_{n \leq x} \frac{\chi(n)\Lambda(n)}{n} = \sum_{d \leq x/m_0} \frac{\chi(d)\mu(d)}{d} \sum_{d_1 \leq x/d} \frac{\chi(d_1) \log d_1}{d_1}. \quad (33)$$

On inserting

$$\sum_{d_1 \leq x/d} \frac{\chi(d_1) \log d_1}{d_1} = -L'(1, \chi) + O\left(\frac{\log(x/d)}{x/d}\right)$$

in this, we obtain

$$\sum_{n \leq x} \frac{\chi(n)\Lambda(n)}{n} = -L'(1, \chi) \sum_{d \leq x/m_0} \frac{\chi(d)\mu(d)}{d} + O\left(\frac{1}{x} \sum_{\substack{d \leq x/m_0 \\ \mu(d)\chi(d) \neq 0}} \log \frac{x}{d}\right). \quad (34)$$

We apply Lemma 8 with $h(n) = \frac{\chi(n)}{n}$ and $f(n) = 1$ together with $\sum_{n \leq x} \chi(n)/n = L(1, \chi) + O(1/x)$ to the latter equation and obtain

$$\begin{aligned} 1 &= \sum_{n \leq x/m_0} \frac{\chi(n)\mu(n)}{n} \left(L(1, \chi) + O\left(\frac{nm_0}{x}\right)\right) \\ &= L(1, \chi) \sum_{n \leq x/m_0} \frac{\chi(n)\mu(n)}{n} + O\left(\frac{m_0}{x} Q_\chi\left(\frac{x}{m_0}\right)\right). \end{aligned}$$

Combining the latter equation with (34) and using the well-known fact that $L(1, \chi) \neq 0$, the result then follows. $\square$

Remark. By using more refined elementary methods [23] one can show that actually, as x tends to infinity,

$$\sum_{n \leq x} \frac{\chi(n)\Lambda(n)}{n} + \frac{L'}{L}(1, \chi) = o(1).$$

Let us consider the case where $\chi = \chi_3$ or $\chi = \chi_4$. Then, for $x > 0$,

$$\left| \sum_{n \leq x} \frac{\chi(n)}{n} - L(1, \chi) \right| \leq \frac{1}{x}, \quad (35)$$

where we use that the non-zero terms in the sum are alternating in sign and monotonically decreasing. The function $\log x/x$ is only decreasing for $x > e$ and a similar argument then shows that, for $x > e$,

$$\left| \sum_{n \leq x} \frac{\chi(n) \log n}{n} + L'(1, \chi) \right| \leq \frac{\log x}{x}. \quad (36)$$

A numerical analysis shows, however, that (36) is still valid for every $x \geq 2$. The implication of these estimates is that for these characters and $x \geq 1$ all the implied constants in the latter lemma and its proof are ≤ 1 . Note that for $x \geq 1$

$$\sum_{\substack{d \leq x \\ \chi(d)\mu(d) \neq 0}} \log \frac{x}{d} = \sum_{\substack{d \leq x \\ \chi(d)\mu(d) \neq 0}} \int_d^x \frac{dt}{t} = \int_1^x \frac{Q_\chi(t)}{t} dt$$

and thus, for $x \geq m_0$,

$$\sum_{\substack{d \leq x/m_0 \\ \chi(d)\mu(d) \neq 0}} \log \frac{x}{d} = \int_1^{x/m_0} \frac{Q_\chi(t)}{t} dt + Q_\chi\left(\frac{x}{m_0}\right) \log m_0.$$

We thus find that, for $x \geq m_0$,

$$x \left| \sum_{n \leq x} \frac{\chi(n) \Lambda(n)}{n} + \frac{L'(1, \chi)}{L(1, \chi)} \right| \leq \left(\frac{L'}{L}(1, \chi) m_0 + \log m_0 \right) Q_\chi\left(\frac{x}{m_0}\right) + \int_1^{x/m_0} \frac{Q_\chi(t)}{t} dt.$$

For $\chi = \chi_3$ we see, using (27), that the right hand side is bounded by

$$\begin{aligned} & \left(\frac{L'}{L}(1, \chi_3) + \frac{\log 2}{2} + \frac{1}{2} \right) \frac{9}{2\pi^2} + \frac{1}{\sqrt{2x}} \left(\frac{L'}{L}(1, \chi_3) + \frac{\log 2}{2} + 1 \right) \\ & + \frac{\log(x/2)}{x} + \frac{1}{x} \left(2 \frac{L'}{L}(1, \chi_3) + \log 2 \right). \end{aligned}$$

For $\chi = \chi_4$ we see, using that $Q_{\chi_4}(t) \leq 4t/\pi^2 + \sqrt{t}/2 + 1$ (Lemma 6), that the right hand side is bounded by

$$\begin{aligned} & \left(\frac{L'}{L}(1, \chi_4) + \frac{\log 3}{3} + \frac{1}{3} \right) \frac{4}{\pi^2} + \frac{1}{\sqrt{3x}} \left(\frac{3}{2} \frac{L'}{L}(1, \chi_4) + \frac{\log 3}{2} + 1 \right) \\ & + \frac{\log(x/3)}{x} + \frac{1}{x} \left(3 \frac{L'}{L}(1, \chi_4) + \log 3 \right). \end{aligned}$$

In the case where $\chi = \chi_3$ it remains to explicitly estimate the latter two sums in (32). We have

$$\sum_{\substack{p^r > \sqrt{x} \\ p \equiv a \pmod{d}}} \frac{\log p}{p^{2r}} \leq \sum_{p^r > \sqrt{x}} \frac{\log p}{p^{2r}} = -\frac{\psi(\sqrt{x})}{x} + 2 \int_{\sqrt{x}}^{\infty} \frac{\psi(t)}{t^3} dt.$$

Using that $0.8t \leq \psi(t) \leq 1.04t$ for $t \geq 17$ (this easily follows from Theorem 10 and Theorem 12 from [27]), we find that

$$\sum_{\substack{p^r > \sqrt{x} \\ p \equiv a \pmod{d}}} \frac{\log p}{p^{2r}} \leq \frac{1.3}{\sqrt{x}} \text{ for } x \geq 289. \quad (37)$$

Furthermore, for every fixed $v > 1$ and every $x > 0$,

$$\frac{\log v}{v-1} \left(1 - \frac{v}{x}\right) \leq \sum_{1 < v^r \leq x} \frac{\log v}{v^r} \leq \frac{\log v}{v-1},$$

where the sum is over the integral powers of v not exceeding x . (These two estimates can also be used in the case where $\chi = \chi_4$.)

Let us define

$$C_+(f) = \sup_{x \geq 1} \left(\sum_{n \leq x} \frac{\Lambda_f(n)}{n} - \tau \log x \right) = B_f + \sup_{x \geq 1} E_f(x),$$

and let $C_-(f)$ be similarly defined, with $\sup$ replaced by $\inf$. Let $\epsilon > 0$ be fixed. Note that the sharpest result the method we followed here allows us to prove, with enough numerical computation, is

$$\begin{aligned} \lim_{x \rightarrow \infty} |E_f(x)| &\leq \left(\frac{L'}{L}(1, \chi_3) + \frac{\log 2}{2} + \frac{1}{2} \right) \frac{9}{4\pi^2} + \epsilon = 0.2769537767 \dots + \epsilon \text{ and} \\ \lim_{x \rightarrow \infty} |E_g(x)| &\leq \left(\frac{L'}{L}(1, \chi_4) + \frac{\log 3}{3} + \frac{1}{3} \right) \frac{2}{\pi^2} + \epsilon = 0.1915268284 \dots + \epsilon, \end{aligned} \quad (38)$$

where $f \in \{g_{3,1}, g_{3,2}\}$ and $g \in \{g_{4,1}, g_{4,3}\}$.

On putting the various effective bounds together we arrive at the following result, after numerical calculations not going beyond the interval $[1, 10^5]$.

Theorem 5 *We have*

- a) $C_-(g_{4,1}) > -1.202$ and $C_+(g_{4,1}) = 0$.
- b) $C_-(g_{4,3}) = \frac{\log 3}{3} - \frac{\log 7}{2} = -0.606750 \dots$ and $C_+(g_{4,3}) = 0$.
- c) $C_-(g_{3,1}) > -1.4$ and $C_+(g_{3,1}) = 0$.
- d) $C_-(g_{3,2}) = -\frac{\log 2}{2} = -0.34657 \dots$ and $C_+(g_{3,2}) < 0.2764$.

On GRH it is much easier to find the C_+ and C_- satisfying (7), which is what will be demonstrated now. By $RH(d)$ we indicate the hypothesis that for every character $\chi \pmod{d}$ every non-trivial zero of $L(s, \chi)$ is on the critical line. Put

$$H(x; d, a) := \sum_{\substack{1 < p^r \leq x \\ p \equiv a \pmod{d}}} \frac{\log p}{p^r} \text{ and } \psi(x; d, a) := \sum_{\substack{n \leq x \\ n \equiv a \pmod{d}}} \Lambda(n).$$

Lemma 10 *For $d \leq 432$ and $(a, d) = 1$, there exists a constant $c_{d,a}$ such that for $x \geq 224$ we have, on $RH(d)$, that*

$$\left| \sum_{\substack{n \leq x \\ n \equiv a \pmod{d}}} \frac{\Lambda(n)}{n} - \frac{\log x}{\varphi(d)} - c_{d,a} \right| \leq \frac{11}{32\pi\sqrt{x}} \{3 \log^2 x + 8 \log x + 16\}, \quad (39)$$

Proof. In [10] it is proved that for $d \leq 432$ and $x \geq 224$ we have, on $\text{RH}(d)$, that

$$|\psi(x; d, a) - \frac{x}{\varphi(d)}| \leq \frac{11}{32\pi} \sqrt{x} \log^2 x. \quad (40)$$

Using the latter estimate and partial integration, the lemma then follows. $\square$

Using the latter lemma we can compute, under GRH, the exact values of $C_-(g_{4,1})$, $C_+(g_{3,1})$ and $C_+(g_{3,2})$.

Theorem 6 *We have*

- a) $C_-(g_{4,1}) = H(197; 4, 1) - \frac{\log(229)}{2} = -0.99076124051235 \dots$, on $\text{RH}(4)$.
- b) $C_-(g_{3,1}) = H(3121; 3, 1) - \frac{\log(3163)}{2} = -1.100304022673 \dots$, on $\text{RH}(3)$.
- c) $C_+(g_{3,2}) = H(5; 3, 2) - \frac{\log 5}{2} = \frac{3}{4} \log 2 - \frac{3}{10} \log 5 = 0.03702 \dots$, on $\text{RH}(3)$.

Proof.

- a) Note that $c_{4,1} = B_{g_{4,1}}$. On applying Lemma 10 with $d = 4$ and $a = 1$, Lemma 4, (37) and using the numerical value for $B_{g_{4,1}}$ given in Section 6, we deduce that $C_-(g_{4,1}) = \min_{q_i \leq 1.79 \cdot 10^9} (H(v_i; 4, 1) - \log(v_{i+1})/2)$, where $5 = v_1 < v_2 < \dots$ are the consecutive prime powers p^r with $p \equiv 1 \pmod{4}$.
- b) In this case we have $C_-(g_{3,1}) = \min_{q_i \leq 2.935 \cdot 10^{10}} (H(q_i; 3, 1) - \log(q_{i+1})/2)$, where $7 = q_1 < q_2 < \dots$ are the consecutive prime powers p^r with $p \equiv 1 \pmod{3}$.
- c) Now $C_+(g_{3,2}) = \max_{w_i \leq 1582079} (H(w_i; 3, 2) - \log(w_i)/2)$, where $2 = w_1 < w_2 < \dots$ are the consecutive prime powers p^r with $p \equiv 2 \pmod{3}$.

9 Connections with Chebyshev's bias for primes

In this section we make some observations that allow us to prove, for example, that $N(x; 3, 2) \geq N(x; 3, 1)$ for every $x \leq x_0$ for some large x_0 , using known numerical observations regarding $\pi(x; 3, 2)$ and $\pi(x; 3, 1)$.

Let $Q_1 = \{q_1, q_2, q_3, \dots\}$ and $Q_2 = \{v_1, v_2, v_3, \dots\}$ be sets of pairwise coprime prime powers that satisfy $q_1 < q_2 < q_3 < \dots$ and $v_1 < v_2 < v_3 < \dots$. Let S_1 denote the set of integers of the form $q_1^{e_1} \dots q_s^{e_s}$ with $q_i \in Q_1$ and $e_i \in \mathbb{Z}_{\geq 0}$ for $1 \leq i \leq s$. Let S_2 be similarly defined, but with Q_1 replaced by Q_2 . Let $\pi_1(x)$, $\pi_2(x)$, count the number of elements in Q_1 , respectively Q_2 , up to x . If $n = q_1^{e_1} \dots q_s^{e_s} \in S_1$, then $m := v_1^{e_1} \dots v_s^{e_s}$ is said to be its *associate* in S_1 . Let $h : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ be a non-increasing function. Put $V_1(x) = \sum_{n \in S_1} h(n)$ and $V_2(x) = \sum_{n \in S_2} h(n)$. In the rest of this section x_0 denotes some arbitrary number.

Lemma 11 *We have*

- a) *If $\pi_1(x) \geq \pi_2(x)$ for $x \geq 0$, then $V_1(x) \geq V_2(x)$ for $x \geq 0$.*
- b) *If $\pi_1(x) \geq \pi_2(x)$ for $x \leq x_0$, then $V_1(x) \geq V_2(x)$ for $x \leq x_0$.*

Proof. a) The assumption implies that if $m \in S_2$, then its associate $n \in S_1$ satisfies $n \leq m$ and $h(n) \geq h(m)$. Thus clearly $V_1(x) \geq V_2(x)$. The proof of part b will be obvious to the reader now. $\square$

Corollary 1 *If $\pi(x; d, a) \geq \pi(x; d, b)$ for $x \leq x_0$, then for $x \leq x_0$ we have both $N(x; d, a) \geq N(x; d, b)$ and $\mu_{g_{d,a}}(x) \geq \mu_{g_{d,b}}(x)$.*

The hypothesis in the corollary is in general not strong enough to infer that $\lambda_{g_{d,a}}(x) \geq \lambda_{g_{d,b}}(x)$ if $x \leq x_0$. However, we have the following easy result.

Lemma 12 *If $M_f(x) \geq M_g(x)$ and $\psi_f(x) \geq \psi_g(x)$ for every $x \leq x_0$, then $\lambda_f(x) \geq \lambda_g(x)$ for $x \leq x_0$.*

Proof. Use (6). □

Corollary 2 *If*

$$\pi(x; d, a) \geq \pi(x; d, b) \text{ and } \sum_{\substack{1 < p^r \leq x \\ p \equiv a \pmod{d}}} \log p \geq \sum_{\substack{1 < p^r \leq x \\ p \equiv b \pmod{d}}} \log p$$

for every $x \leq x_0$, then $\lambda_{g_{d,a}}(x) \geq \lambda_{g_{d,b}}(x)$ for $x \leq x_0$.

In the proof of Theorem 7 we will put Corollary 2 a few times to action.

10 The proof of Theorem 2

The proof of Theorem 2 will easily follow from the following theorem.

Theorem 7 *For every x we have $\lambda_{g_{3,2}}(x) \geq \lambda_{g_{3,1}}(x)$, $\lambda_{g_{3,2}}(x) \geq \lambda_{g_{4,1}}(x)$ and $\lambda_{g_{4,3}}(x) \geq \lambda_{g_{3,1}}(x)$. For $x \geq 7$ we have $\lambda_{g_{4,3}}(x) \geq \lambda_{g_{4,1}}(x)$.*

Note that

$$e^{\lambda_{g_{d,a}}(x)} = \prod_{\substack{n \leq x \\ p|n \Rightarrow p \equiv a \pmod{d}}} n.$$

In the proof of Theorem 7 we will make use of the following lemma.

Lemma 13 *We have $\psi_{g_{3,1}}(x) \leq 0.50456x$ for $x \geq 0$, $\psi_{g_{3,2}}(x) \geq 0.335x$ for $x \geq 5$, $\psi_{g_{4,1}}(x) \leq 0.50456x$ for $x \geq 0$ and $\psi_{g_{4,3}}(x) \geq 0.48508x$ for $x \geq 127$.*

Proof. Let $d \leq 13$ and $(a, d) = 1$. Then $|\psi(x; d, a) - x/\varphi(d)| \leq \sqrt{x}$ for $224 \leq x \leq 10^{10}$ by [26, Theorem 1] and $|\psi(x; d, a) - \frac{x}{\varphi(d)}| < 0.004560 \frac{x}{\varphi(d)}$ for $x \geq 10^{10}$ by [26, Theorem 5.2.1]. From these inequalities the lemma follows after some computation. □

In our proof we consider inequalities of the form

$$\log^\tau\left(\frac{x}{r}\right) \frac{\left(1 - \frac{C_+}{\log(x/r)}\right)^{\tau+1}}{\left(1 - \frac{C_-}{\log(x/r)}\right)} \geq c_1 \log^\tau\left(\frac{x}{s}\right) \frac{\left(1 - \frac{C'_-}{\log(x/s)}\right)^{\tau+1}}{\left(1 - \frac{C'_+}{\log(x/s)}\right)}, \quad (41)$$

where all variables and constants are real numbers with τ, r, s and c_1 positive, $C_- \leq C_+$, $C'_- \leq C'_+$ and $x \geq x_0 := \max\{\exp(C'_+)s, \exp(C_+)r\}$. This inequality can be rewritten as

$$1 + \frac{C'_- - C_+ + \log(s/r)}{\log(x/s) - C'_-} \geq \left[c_1 \frac{\left(1 + \frac{C'_+ - C'_-}{\log(x/s) - C'_+}\right)^{\frac{1}{\tau}}}{1 + \frac{C_- - C_+}{\log(x/r) - C_-}} \right]^{\frac{1}{\tau}}. \quad (42)$$

Note that for $x > x_0$ the right hand side is a non-increasing function of x . If $C'_- + \log s \leq C_+ + \log r$, the left hand side is non-decreasing, whereas if the latter inequality is not satisfied the left hand side asymptotically decreases to 1. We thus arrive at the following conclusion.

Lemma 14 *If $\log s + C'_- \leq C_+ + \log r$ and (41) is satisfied for some $x_1 > x_0$, then (41) is satisfied for every $x \geq x_1$. If $\log s + C'_- > C_+ + \log r$, and the right hand side of (42) does not exceed 1 for some $x_1 > x_0$, then (41) is satisfied for every $x \geq x_1$.*

Proof of Theorem 7.

$\lambda_{g_{3,2}}(x)$ versus $\lambda_{g_{3,1}}(x)$. Using Lemma 13 we infer that

$$\lambda_{g_{3,2}}(x) \geq \sum_{n \leq \frac{x}{5}} g_{3,2}(n) \psi_{g_{3,2}}\left(\frac{x}{n}\right) \geq 0.335 \mu_{g_{3,2}}\left(\frac{x}{5}\right), \text{ and that}$$

$$\lambda_{g_{3,1}}(x) = \sum_{n \leq x} g_{3,1}(n) \psi_{g_{3,1}}\left(\frac{x}{n}\right) = \sum_{n \leq \frac{x}{7}} g_{3,1}(n) \psi_{g_{3,1}}\left(\frac{x}{n}\right) \leq 0.50456 \mu_{g_{3,1}}\left(\frac{x}{7}\right).$$

With $d = 3$, $a = 2$ and $b = 1$ the conditions of Corollary 2 are satisfied for every $x < 196699$ (but not for $x = 196699$ as $\psi_{g_{3,1}}(196699) > \psi_{g_{3,2}}(196699)$). Thus we certainly may assume that $x > 1900$. Using the estimates $C_{3,1} < 0.302$ and $C_{3,2} > 0.703$ we then deduce from Lemma 1, Theorem 5 and Lemma 14 that $0.335 \mu_{g_{3,2}}(x/7) > 0.50456 \mu_{g_{3,1}}(x/7)$.

$\lambda_{g_{3,2}}(x)$ versus $\lambda_{g_{4,1}}(x)$. The conditions of Corollary 2 are now satisfied for every $x \leq 10^7$ (the smallest x for which the conditions are not satisfied is not known, but must be less than 10^{82} by [12]). Thus we certainly may assume that $x > 4600$. Then reasoning as before we infer that $\lambda_{g_{3,2}}(x) \geq 0.335 \mu_{g_{3,2}}(x/5) \geq 0.50456 \mu_{g_{4,1}}(x/5) \geq \lambda_{g_{4,1}}(x)$.

$\lambda_{g_{3,2}}(x)$ versus $\lambda_{g_{4,1}}(x)$. The conditions of Corollary 2 are now satisfied for every $x \leq 10^7$ (the smallest x for which the conditions are not satisfied is not known, but must be less than 10^{82} by [12]). Thus we may assume that $x > 199000$. Then it is seen that $\lambda_{g_{4,3}}(x) \geq 0.4594 \mu_{g_{4,3}}(x/59) \geq 0.50456 \mu_{g_{3,1}}(x/5) \geq \lambda_{g_{3,1}}(x)$.

$\lambda_{g_{4,3}}(x)$ versus $\lambda_{g_{4,1}}(x)$. For $7 \leq x \leq 1.1 * 10^6$ one directly verifies the inequality (note that Corollary 2 cannot be used this time). For $x > 1.1 * 10^6$ one deduces, proceeding as before, that $\lambda_{g_{4,3}}(x) \geq 0.48508 \mu_{g_{4,3}}(x/127) \geq 0.50456 \mu_{g_{4,1}}(x/5) \geq \lambda_{g_{4,1}}(x)$.

It remains to establish Theorem 2.

Proof of Theorem 2. We only deal with $N(x; 4, 3)$ versus $N(x; 4, 1)$, the other cases following at once from Theorem 7 and (21). Put $\delta(x) = \lambda_{g_{4,3}}(x) - \lambda_{g_{4,1}}(x)$. By Theorem 7 we have $\delta(x) \geq 0$ for $x \geq 7$. Using this and (21) we infer that

$$\begin{aligned} N(x; 4, 3) - N(x; 4, 1) &= \frac{\delta(x)}{\log x} + \int_2^7 \frac{\delta(t) dt}{t \log^2 t} + \int_7^x \frac{\delta(t) dt}{t \log^2 t} \\ &\geq \int_2^7 \frac{\delta(t) dt}{t \log^2 t} = \frac{\log 5 - \log 3}{\log 7} > 0, \end{aligned}$$

for $x \geq 7$. For $x < 7$ the result is clearly true. □

Acknowledgement. T. Dokshitzer was so kind as to redo some of the computations (carried out in Maple and Quickbasic) in PARI. K. Ford, R. Hudson and M. Rubinstein helpfully provided me with some numerical data regarding Chebyshev's bias (and with a preprint of [12]). I'd like to thank O. Ramaré for pointing out reference [10]. He also informed me that he is working on a paper [25] in which he is able to roughly half the values given in (38) (which is not enough to give an unconditional proof of any of the claims in Theorem 6). H. te Riele kindly did the computations required to validate parts a and b of Theorem 6, taking respectively 20 and 5 minutes CPU time on a 300MHz SGI processor.

References

- [1] C. Bays and R.H. Hudson, Details of the first region of integers x with $\pi_{3,2}(x) < \pi_{3,1}(x)$, *Math. Comp.* **32** (1978), 571-576.
- [2] P.L. Chebyshev, Lettre de M. le Professeur Tchébychev à M. Fuss sur un nouveau théorème relatif aux nombres premiers contenus dans les formes $4n+1$ et $4n+3$, *Bull. Classe Phys. Acad. Imp. Sci. St. Petersburg* **11** (1853), 208.
- [3] H. Cohen, *Advanced topics in computational number theory*, GTM **193**, Springer-Verlag, New York, 2000.
- [4] H. Cohen, High precision computation of Hardy-Littlewood constants, draft of a preprint, available at <http://www.math.u-bordeaux/~cohen/>.
- [5] H. Cohen and F. Dress, Estimations numériques du reste de la fonction sommatoire relative aux entiers sans facteur carré, *Publ. Math. Orsay* 88/02 (1988), 73-76.
- [6] J.H. Conway and N.J.A. Sloane, Sphere packings, lattices and groups, Third edition, Springer-Verlag, New York, 1999.
- [7] D.A. Cox, The arithmetic-geometric mean of Gauss, *Enseign. Math.* **30** (1984), 275-330.
- [8] H. Davenport, *Multiplicative number theory*, Third revised edition, Springer-Verlag, New York, 2000.
- [9] K. Dilcher, Generalized Euler constants for arithmetical progressions, *Math. Comp.* **59** (1992), 259-282.
- [10] P. Dusart, Autour de la fonction qui compte le nombre de nombres premiers, PhD thesis, Université de Limoges, 1998.
- [11] S. Finch, Mathematical constant web pages, <http://www.mathsoft.com/asolve/constant/constant.html>

- [12] K. Ford and R.H. Hudson, Sign changes in $\pi_{q,a}(x) - \pi_{q,b}(x)$, submitted for publication.
- [13] E. Landau, Über die zu einem algebraischen Zahlkörper gehörige Zetafunktion und die Ausdehnung der Tschebyscheffschen Primzahlentheorie aus das Problem der Verteilung der Primideale, *J. reine angew. Math.* **125** (1903), 64-188.
- [14] E. Landau, Über die Einteilung der positiven ganzen Zahlen in vier Klassen nach der mindest Anzahl der zu ihrer additiven Zusammensetzung erforderlichen Quadrate, *Arch. der Math. und Phys.* **13** (1908), 305-312.
- [15] J.E. Littlewood, Distribution des nombres premiers, *C.R. Acad. Sci. Paris* **158** (1914), 1869-1872.
- [16] G. Martin, Asymmetries in the Shanks-Rényi Prime Number Race, proceedings of the Millennial Conference on Number Theory (Urbana, IL), to appear.
- [17] S.D. Miller and G. Moore, Landau-Siegel zeroes and black hole entropy, *Asian J. Math.* **4** (2000), 183-211.
- [18] P. Moree, Approximation of singular series and automata, *Manuscripta Math.* **101** (2000), 385-399.
- [19] P. Moree and J. Cazarán, On a claim of Ramanujan in his first letter to Hardy, *Exposition. Math.* **17** (1999), 289-311.
- [20] P. Moree and H.J.J. te Riele, The hexagonal versus the square lattice, in preparation.
- [21] L. Moser and R.A. MacLeod, The error term for the squarefree integers, *Canad. Math. Bull.* **9** (1966), 303-306.
- [22] W. Narkiewicz, *The development of prime number theory. From Euclid to Hardy and Littlewood*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2000.
- [23] V. Nevanlinna, On constants connected with the prime number theorem for arithmetic progressions, *Ann. Acad. Sci. Fenn. Ser. A I* **539** (1973), 11 pp..
- [24] A.G. Postnikov, *Introduction to analytic number theory*, AMS translations of mathematical monographs 68, AMS, Providence, Rhode Island, 1988.
- [25] O. Ramaré, Sur un théorème de Mertens, in preparation.
- [26] O. Ramaré and R. Rumely, Primes in arithmetic progressions, *Math. Comp.* **65** (1996), 397-425.
- [27] J.B. Rosser and L. Schoenfeld, Approximate formulas for some functions of prime numbers, *Illinois Journal Math.* **6** (1962), 64-94.

- [28] M. Rubinstein and P. Sarnak, Chebyshev's bias, *Experiment. Math.* **3** (1994), 173-197.
- [29] P. Schmutz Schaller, Geometry of Riemann surfaces based on closed geodesics, *Bull. Amer. Math. Soc. (N.S.)* **35** (1998), 193-214.
- [30] L. Schoenfeld, Sharper bounds for the Chebyshev functions $\theta(x)$ and $\psi(x)$. II, *Math. Comp.* **30** (1976), 337-360.
- [31] J.-P. Serre, Divisibilité de certaines fonctions arithmétiques, *Enseign. Math.* **22** (1976), 227-260.
- [32] D. Shanks, The second-order term in the asymptotic expansion of $B(x)$, *Math. Comp.* **18** (1964), 75-86.
- [33] J.M. Song, Sums over nonnegative multiplicative functions over integers without large prime factors. I, *Acta Arith.* **97**, 329-351.
- [34] P. Turán, *On a new method of analysis and its applications*, John Wiley and Sons, Inc., New York, 1984.
- [35] P. Turán, Collected papers of Paul Turán, Vol. 1-3, Ed. P. Erdős. Akadémiai Kiadó, Budapest, 1990.
- [36] K.S. Williams, Mertens' theorem for arithmetic progressions, *J. Number Theory* **6** (1974), 353-359.
- [37] A. Wintner, On the distribution of the remainder term of the prime number theorem, *Amer. J. Math.* **63** (1941), 233-248.
- [38] E. Wirsing, Das asymptotische Verhalten von Summen über multiplikative Funktionen, *Math. Ann.* **143** (1961), 75-102.